arXiv:1212.6092v1 [math.CO] 25 Dec 2012

A NOTE ON STRONG EDGE-COLORINGS OF 2-DEGENERATE GRAPHS

RONG LUO† AND GEXIN YU‡

Abstract. We show that every 2-degenerate graph with maximum degree Δ has a strong edge-coloring with at most $8\Delta - 4$ colors.

1. Introduction

A *strong edge-coloring* of a graph G is an edge-coloring in which every color class is an induced matching; that is, no edge can be incident to two edges with the same color. The *strong chromatic index* $\chi'_s(G)$ is the minimum number of colors in a strong edge-coloring of G. This notion was introduced by Fouquet and Jolivet (1983, [6]) and one of the main open problems was proposed by Erdős and Nešetřil during a seminar in Prague in 1985, as follows,

Conjecture 1.1 (Erdős and Nešetřil, 1985)**.** *If G is a simple graph with maximum degree Δ, then $\chi'_s(G) \le 5\Delta^2/4$ if Δ is even, and $\chi'_s(G) \le (5\Delta^2 - 2\Delta + 1)/4$ if Δ is odd.*

This conjecture is true for $\Delta \le 3$, see [1, 9]. For $\Delta = 4$, Cranston [4] showed that $\chi'_s(G) \le 22$, two more than the conjectured upper bound 20. Chung, Gyárfás, Trotter, and Tuza (1990, [3]) confirmed the conjecture for $2K_2$-free graphs. Using probabilistic methods, Molloy and Reed [10] proved that for sufficiently large Δ, every graph with maximum degree Δ has strong chromatic index at most $1.998\Delta^2$. Readers are referred to Problem 17 in Chapter 6 of [11] for more information about strong edge colorings of graphs.

Sparse graphs have also attracted a lot of attention. Faudree, Gyárfás, Schelp, and Tuza (1989, [5]) showed that every planar graph with maximum degree Δ has strong chromatic index at most $4\Delta + 4$. Hocquarda, Ochemb, and Valicov [7] showed a tight result which claims that every non-trivial outerplanar graph can be strongly edge-colored with at most $3\Delta - 3$ colours. Hocquard, Montassier, Raspaud, and Valicov [8] gave the optimal upper bound on χ'_s for subcubic graphs with certain maximum average degrees.

A graph is 2-*degenerate* if every subgraph has minimum degree at most two. Outplanar graphs, non-regular subcubic graphs, and planar graphs with girth at least six are all 2-degenerate graphs. Chang and Narayanan (2012, [2]) recently proved that a 2-degenerate graph with maximum degree Δ has strong chromatic index at most $10\Delta - 10$. They actually proved the following stronger statement (see paragraphs 4 and 5 in [2]).

Date: December 27, 2012.

† The author's research is supported in part by a NSF-China grant: NSFC 11171228.

‡ The author's research is supported in part by a NSA grant.

Theorem 1.2. *(Chang and Narayanan, [2]) Let G be a 2-degenerate graph with maximum degree $\Delta > 0$. Let $B = \{1, 2, \cdots, 5\Delta - 5\}$ and $B' = \{1', 2', \cdots, (5\Delta - 5)'\}$. Then G has a strong edge-coloring with colors from $B \cup B'$ so that*
(a) Every pendant edge (if any) is colored with a color in B;
(b) If a pendant edge is colored with a color $c \in B$, then no edge within distance 1 to the edge is colored with c'.

In this note, we prove a stronger result with a better upper bound. The induction in our proof is similar to their proof, except that we add an extra condition (3). This extra condition helps fix a gap in their proof (in the third paragraph on Page 5 in [2], if some edge incident to v has color c', then the priming process fails as one cannot color ww' with c'). Also in our proof, instead of considering a subgraph by deleting one edge from G, we delete a bunch of edges and try to color those deleted edges one by one, and this helps us improve the upper bound.

Theorem 1.3. *Let G be a 2-degenerate graph with maximum degree at most Δ. Let $B = \{1, 2, \cdots, 4\Delta - 2\}$ and $B' = \{1', 2', \cdots, (4\Delta - 2)'\}$. Then G has a strong edge-coloring with colors from $B \cup B'$ so that*
(1) Every pendant edge is colored with a color in B;
(2) If a pendant edge is colored with a color $c \in B$, then no edge within distance 1 to the edge is colored with c';
(3) No pair of colors $\{c, c'\}$ appears at the same vertex.

As a corollary, every 2-degenerate graph with maximum degree Δ has strong chromatic index at most $8\Delta - 4$.

2. Proof of Theorem 1.3

In this section, we prove Theorem 1.3. We will use the following notion. A k-vertex (k^--vertex, k^+-vertices) is a vertex of degree k (at most k, at least k). If a k-vertex (k^--vertex, k^+-vertex) u is a neighbor of vertex v, then we also call u to be a k-neighbor (k^--neighbor, k^+-neighbor) of v. Let $B = \{1, 2, \cdots, 4\Delta - 2\}$ and $B' = \{1', 2', \cdots, (4\Delta - 2)'\}$ be the sets of colors. For a color $c \in B \cup B'$, denote c' to be the corresponding color in the other color set (thus $(c')' = c$). For $A \subseteq B$ or $A \subseteq B'$, let $A' = \{c' : c \in A\}$. If f is a strong edge coloring of G (or its subgraph), then we use $C_f(v)$ to denote the set of colors of the edges incident with $v \in V(G)$.

First of all, we quote the following lemma on 2-degenerated graphs.

Lemma 2.1. *(Lemma 3, [2]) Let G be a 2-degenerate graph. Then G has a vertex v with degree at least 3 that has at most two 3^+-neighbors.*

Proposition 2.2. *Let uv be a pendant edge in G with $d(u) = 1$. If f is an edge-coloring of G with properties in Theorem 1.3 so that $f(uv) = c$, then the coloring g, which differs from f only on uv with $g(uv) = c'$, is a strong edge-coloring which satisfies (2) and (3) but violates (1) (only) at the pendant edge uv.*

Proof. It is clear that uv is the only pendant edge colored with a color in B' in g. Since $f(uv) = c$, by (2), $c, c' \notin \cup_{u\in N(v)} C_f(u) = \cup_{u\in N(v)} C_g(u)$ and c, c' do not appear on the edges incident with v (except uv) in f thus also in g. So g satisfies (2) and (3). $\square$

Proof of Theorem 1.3. Let G be a smallest counterexample to Theorem 1.3. That is, G is a 2-degenerate graph with maximum degree at most Δ and very proper subgraph of G has a desired strong edge coloring.

By Lemma 2.1, let v be a vertex adjacent to at most two 3^+-vertices. Let $d = d(v) - 2$, and $v_1, \cdots, v_d$ be 2^--neighbors of v, and u and w be the other two neighbors of v. We have the following claim.

Claim 2.3. *$d(v_i) = 2$ for $i = 1, \cdots d$.*

Proof. Without loss of generality, suppose that v_1 is a 1-neighbor of v. Let f be a strong edge-coloring of $G - v_1$ satisfying (1)-(3). Let $A = \cup_{x\in N(v)} C_f(x)$. Then $(A \cap B')' \subseteq B$ and $(A \cap B) \cup (A \cap B')' \subseteq B$, and

$$|(A \cap B) \cup (A \cap B')'| \le |A| \le \sum_{x\in N(v)} d(x) - 1 \le 2\Delta + 2(d(v) - 3) - 1 \le 4\Delta - 7.$$

Therefore let $f(vv_1) = c$, where $c \in B - ((A\cap B)\cup(A\cap B')')$. As $c, c' \notin A$, f is a strong edge-coloring of G.

To see a contradiction, we now show that f satisfies (1)-(3). Since $f(vv_1) = c \in B$, Property (1) is met. Since $c \notin A \cap B$ and $c \notin (A \cap B')'$, $c \notin A \cap B$ and $c' \notin A \cap B'$, that is, $c \notin A$. Therefore, Properties (2) and (3) are met. $\square$

By Claim 2.3, for each $i = 1, 2, \cdots, d$, let $N(v_i) = \{v, u_i\}$. Consider $G_1 = G - \{vv_i : 1 \le i \le d\}$. Then G_1 has a desired strong edge-coloring f. As v_iu_i is a pendant edge in G_1, it is colored with a color in B.

We now consider a new edge-coloring g of G which may only differ from f on edges v_iu_i. For each edge v_iu_i, if $(f(v_iu_i))' \notin \{f(vu), f(vw)\}$, then let $g(v_iu_i) = (f(v_iu_i))'$, otherwise, $g(v_iu_i) = f(v_iu_i)$. By Proposition 2.2, g is a strong edge-coloring of G_1 which satisfies (2) and (3). We are going to color the edge vv_i one by one with available colors in B. As the edges v_iu_i are not pendent edges in G, (1) is not violated in the new coloring. Note that the more edges vv_i are colored, the less choices for the remaining uncolored edges, so we only need to consider when only one edge (say vv_1) is yet to color. We still use g to denote the strong edge-coloring of $G - vv_1$ which is an extension of the coloring g of G_1.

Let $S = \{g(vu), g(vw), g(v_1u_1)\} \cap B'$. Then $S' \subseteq B$. Note that by the construction of g, $g(vv_i) \in B$ for each $i = 2, 3, \cdots, d$ and furthermore $g(v_iu_i) = c \in B$ if and only if $c' \in S$, thus if and only if $c \in S'$. Let

$$\Omega = [\{g(vu), g(vw)\} \cap B] \cup S' \cup \{g(vv_i) : 2 \le i \le d\} \cup ([C_g(u_1) \cup C_g(u) \cup C_g(w)] \cap B).$$

Since

$$|\Omega| \le (d(u) + d(w) + (d(v) - 3) + d(u_1) \le 4\Delta - 3,$$

pick one color $t \in B \setminus \Omega$ and let $g(vv_1) = t$, so we obtain an edge-coloring of G. By the choice of t, $g(vv_1) = t \neq g(v_1u_1)$ and no edge within distance 1 to vv_1 is colored with t except those on v_iu_i with $i \geq 2$. But if $v_iu_i = t$ for some $i \geq 2$, then $t' \in \{f(vu), f(vw)\}$, so $t \in \Omega$, a contradiction. Therefore g is a strong edge-coloring of G.

Lastly, we show that g satisfies Properties (1)-(3). Since none of v_1u_1 and vv_1 are pendant edges in G, (1) and (2) are satisfied. So we just need to show that $g(v_1u_1) \neq t'$ and $t' \notin C_g(v)$. If there is an edge incident with v_1 or v is colored with t', then $e \in \{v_1u_1, vw, vu\}$ since $g(vv_i) \in B$ for each $i = 2, 3 \cdots$. Hence $f(e) = t' \in S$ and thus $t = S' \subseteq \Omega$, a contradiction. Therefore (3) is met. This finishes the proof of Theorem 1.3.

3. Final remarks

In [2], they also studied the strong chromatic index of a chordless graph G and showed that $\chi'_s(G) \leq 8\Delta - 8$. Their proof is very similar to the one for 2-degenerate graphs, thus also has a similar gap. We can improve their result to the following

Theorem 3.1. *Let G be a chordless graph with maximum degree at most Δ. Let $B = \{1, 2, \cdots, 3\Delta - 1\}$ and $B' = \{1', 2', \cdots, (3\Delta - 1)'\}$. Then G has a strong edge-coloring with colors from $B \cup B'$ so that*
(1) Every pendant edge is colored with a color in B;
(2) If a pendant edge is colored with a color $c \in B$, then no edge within distance 1 to the edge is colored with c';
(3) No pair of colors $\{c, c'\}$ appears at the same vertex.

So if a chordless G has maximum degree Δ, then $\chi'_s(G) \leq 6\Delta - 2$. The proof is very similar to that of Theorem 1.3, so we omit it here.

We do not believe that the bounds in this note are optimal. Let G be the graph consisting of a triangle with $D - 2$ leaves on each of the vertices on the triangle. Then $\Delta(G) = D$ and $\chi'_s(G) = 3\Delta - 3$, but G is 2-degenerate and chordless. So $\chi'_s(G) \geq 3\Delta - 3$ for any 2-degenerate or chordless graph G with maximum degree Δ.

References

[1] L. D. Andersen, The strong chromatic index of a cubic graph is at most 10. Topological, algebraical and combinatorial structures. Frolík's memorial volume. *Discrete Math.* 108 (1992), no. 1-3, 231–252.

[2] G.J. Chang and N. Narayanan, Strong Chromatic Index of 2-Degenerate Graphs, *J. Graph Theory*, DOI: 10.1002/jgt.21646.

[3] F. R. K. Chung, A. Gyárfás, W. T. Trotter, and Z. Tuza, The maximum number of edges in $2K_2$-free graphs of bounded degree. *Discrete Mathematics*, 81(2):129–135, 1990.

[4] D. W. Cranston, Strong edge-coloring of graphs with maximum degree 4 using 22 colors. *Discrete Math.* 306 (2006), no. 21, 2772–2778.

[5] R. J. Faudree, A. Gyárfás, R. H. Schelp, and Zs. Tuza, Induced matchings in bipartite graphs. *Discrete Math.* 78 (1989), no. 1-2, 83–87.

[6] J. L. Fouquet and J. L. Jolivet, Strong edge-coloring of graphs and applications to multi-k-gons. *Ars Combinatoria*, 16A:141–150, 1983.

[7] H. Hocquard, P. Ochem, and P. Valicov, Strong edge colouring and induced matchings. Submitted.

[8] H. Hocquard, M. Montassier, A. Raspaud, and P. Valicov, On strong edge-colouring of graphs. Submitted.

[9] P. Horák, Q. He, W. T. Trotter, Induced matchings in cubic graphs. *J. Graph Theory* 17 (1993), no. 2, 151–160.

[10] M. Molloy, B. Reed, A bound on the strong chromatic index of a graph. *J. Combin. Theory Ser. B* 69 (1997), no. 2, 103–109.

[11] M. Steibitz, D. Scheide, B. Toft, and L. M. Favrholdt, Graph Edge Coloring, A John Wiley & Sons, Inc, 2012.

† Department of Mathematics, West Virginia University, Morgantown, WV 26506-6310.
E-mail address: `rluo@math.wvu.edu`

‡ Department of Mathematics, College of William and Mary, Williamsburg, VA, 23185.
E-mail address: `gyu@wm.edu`